

X-RAY SPECTROSCOPIC EVIDENCE FOR INTERMEDIATE MASS BLACK HOLES: COOL ACCRETION DISKS IN TWO ULTRA-LUMINOUS X-RAY SOURCES

J. M. MILLER^{1,4}, G. FABBIANO¹, M. C. MILLER², A. C. FABIAN³

Subject headings: Black hole physics – X-rays:stars

Draft version February 2, 2008

ABSTRACT

We have analyzed an *XMM-Newton* observation of the nearby spiral galaxy NGC 1313, which contains two “ultra-luminous” X-ray (ULX) sources. We measure isotropic luminosities of $L_X = 2.0 \times 10^{40} \text{ erg s}^{-1}$ and $L_X = 6.6 \times 10^{39} \text{ erg s}^{-1}$ for NGC 1313 X-1 and X-2 (0.2–10.0 keV, assuming a distance of 3.7 Mpc). The spectra *statistically require* soft and hard spectral components to describe the continuum emission; some prior studies of ULXs have claimed cool soft components with lower statistics. The improvement over several single-component models exceeds the 8σ level of confidence for X-1; the improvement for X-2 is significant at the 3σ level. The soft components in these ULX spectra are well-fit by multi-color disk blackbody models with color temperatures of $kT \simeq 150 \text{ eV}$. This temperature differs markedly from those commonly measured in the spectra of stellar-mass ($10 M_\odot$) black holes in their brightest states ($kT \simeq 1 \text{ keV}$). It is expected that the temperature of an accretion disk orbiting a black hole should decrease with increasing black hole mass. If the soft components we measure are due to emission from the inner region of an accretion disk, and disks extend close to the innermost stable circular orbit at the accretion rates being probed, the low color temperatures may be interpreted as spectroscopic evidence of black holes with intermediate masses: $M_{BH} \simeq 10^3 M_\odot$. Simple Eddington scaling arguments suggest a minimum mass of $M_{BH} \sim 10^2 M_\odot$. NGC 1313 X-1 and X-2 are found in optical nebulae (Pakull & Mirioni 2002), which may indicate that anisotropic emission geometries are unlikely to account for the fluxes observed.

1. INTRODUCTION

ULX sources may be defined as point-like off-nuclear X-ray sources in normal galaxies for which measured luminosities exceed the isotropic Eddington limit for a stellar-mass (less than approximately $10 M_\odot$) black hole. The existence of such sources was first revealed with *Einstein* (Fabbiano 1989). Variability has been observed in many ULX sources on the timescales of months and years (in rare cases, on shorter timescales), which suggests that they are accreting objects.

Although black holes with intermediate mass (IMBHs; $10^{2-5} M_\odot$) provide an attractive explanation for the nature of ULX sources, strong evidence for this interpretation has been lacking — especially from X-ray spectroscopic studies. Anisotropic emission from stellar-mass black holes may be able to account for the flux observed in some ULX sources (King et al. 2001). Single-component fits with the multi-color disk blackbody model (MCD; Mitsuda et al. 1984) have measured color temperatures above those commonly reported in stellar-mass black holes ($kT = 1\text{--}2 \text{ keV}$; see, e.g., Sobczak et al. 2000, Makishima et al. 2000); as inner disk temperatures should fall with increasing black hole mass, these findings again point towards stellar-mass black holes or Kerr black holes.

NGC 1313 is a nearby spiral galaxy ($d = 3.7 \text{ Mpc}$, Tully 1988). Two ULX sources — NGC 1313 X-1 X-2 — are associated with this galaxy; approximately 1 kpc and 8 kpc from the photometric center of the galaxy, respectively (Colbert et al. 1995). Variability was observed in *ROSAT*/HRI observations of these sources (Colbert & Ptak 2002). Colbert and

Mushotzky (1999) claim evidence for a cool ($kT = 120 \text{ eV}$) disk in fits to an *ASCA* spectrum of NGC 1313 X-1 with a model consisting of MCD and power-law components; however the spectrum is fit acceptably by a simple power-law ($\Gamma = 1.74$, $\chi^2/\text{d.o.f.} = 244/259$). Similarly, a cool disk may be implied in joint fits to *BeppoSAX*/MECS and *ROSAT*/PSPC spectra of M81 X-9 (La Parola et al. 2001) but not in fits to MECS and LECS spectra (both aboard *BeppoSAX*).

We have analyzed the archival *XMM-Newton* spectra of NGC 1313 X-1 and X-2. Separate soft and hard components are statistically required to describe the spectra. We measure low temperatures ($kT \simeq 150 \text{ eV}$) for the soft components. We interpret the soft component as arising from the inner region of an accretion disk, and explore the implications of cool accretion disks for the mass of black holes, if such objects power X-1 and X-2.

2. DATA REDUCTION AND ANALYSIS

NGC 1313 was observed by *XMM-Newton* on 17 October 2000 starting at 03:59:23 (UT). We used only the EPIC data for this analysis. The EPIC cameras were operated in “PrimeFull-Window” mode with the “medium” optical blocking filter. The *XMM-Newton* reduction and analysis suite SAS version 5.3.3 was used to filter the standard pipeline event lists, to detect sources within the field, and to make spectra and responses. The pipeline processing and our own both failed to produce an event list for the pn camera. We therefore restricted our analysis to the MOS-1 and MOS-2 cameras. Application of the standard time filtering resulted in a net exposure of 29.3 ksec.

¹Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, jmmiller@head-cfa.harvard.edu

²Department of Astronomy, University of Maryland, College Park, MD, 20742

³Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, United Kingdom

⁴National Science Foundation Astronomy and Astrophysics Fellow

The source locations were determined by running the SAS tool “edetect_chain”. With this tool, we find NGC 1313 X-1 at 3h18m19.99s, $-66^{\circ}29'10.97''$, and NGC 1313 X-2 at 3h18m22.34s, $-66^{\circ}36'03.68''$ (J2000). The tool returns an error of 0.2” for these positions; an uncertainty of 4” may be more appropriate (e.g., Foschini et al. 2002). Source counts were extracted in a circle within 24” of the detected source position. Background counts were extracted in an annulus between 24”–30”. To create spectra, we then applied the selection criteria described in the MPE “cookbook.” These selections are as follows: we set “FLAG=0” to reject events from bad pixels and events too close to the CCD chip edges, event patterns 0–12 were allowed, and the MOS spectral channels were grouped by a factor of 15. Response files were made using the SAS tools “rmfgen” and “arfgen.” Spectral files were grouped to require at least 20 counts per bin before fitting to ensure the validity of χ^2 statistics.

Model spectra were fit to the data using XSPEC version 11.2 (Arnaud 1996). The MOS-1 and MOS-2 spectra were fit jointly with an overall normalizing constant. The constant indicates that the overall flux normalizations of these cameras differ by less than 5%. Models were fit to the spectra in the 0.2–10.0 keV band. Systematic errors were not added to the spectra. Errors quoted in this work are at the 90% confidence level. Using the SAS tool “epatplot” we found photon pile-up to be negligible in our source spectra; this was confirmed with the HEASARC tool “PIMMS” using the parameters reported by Colbert & Mushotzky (1999) for X-1 and X-2.

3. RESULTS

Lightcurves of the source event lists do not show strong variability on the timescale of this observation. We therefore proceeded to make fits to the time-averaged spectra. The results of joint fits to the MOS-1 and MOS-2 spectra in the 0.2–10.0 keV band are listed in Table 1.

We began by fitting single-component models commonly applied to ULX sources, modified by photoelectric absorption (via the “phabs” model within XSPEC). For both X-1 and X-2, MCD, thermal Bremsstrahlung, and Raymond-Smith plasma models all fail to yield acceptable fits. Simple power-law models provide improved but statistically unacceptable fits. Broken power-law models with breaks in the 0.2–10.0 keV range also yield improved but statistically unacceptable fits.

An observation of NGC 1313 with *Chandra* finds that the diffuse emission in our *XMM-Newton* extraction region is not well-described with any thermal model; the best model is a $\Gamma = 1.7$ power-law with an unabsorbed flux of $F_{0.3-7.0} = 5.9 \times 10^{-14}$ erg/s (A. Kong, priv. comm.). This signals that the soft components may indeed arise in accretion disks, and we explored fits with a model consisting of MCD and power-law components, and the “CompTT” model. The MCD model describes a standard Shakura-Sunyaev (1973) accretion disk as a series of blackbody annuli. The latter model describes Compton-upscattering in a corona of optical depth τ with electron temperature kT_{corona} from a Wien distribution of soft seed photons (Titarchuk 1994); we used the version of the model which assumes a disk geometry for the seed photons. The coronal temperature could not be constrained and we therefore fixed $kT_{\text{corona}} = 50$ keV, which is a moderate value. X-2 is fit acceptably by both of these models; the MCD plus power-law model is a significantly better fit to the spectrum of NGC X-1. The inability of single-component models to describe the spectra




FIG. 1.— The data/model ratios from fits to the MOS-1 (black) and MOS-2 (red) spectra of NGC 1313 X-1 with standard ULX models (the spectra are rebinned for visual clarity). The ratios correspond to fits with: a simple power-law, the multicolor disk blackbody model, and a model with both components (see Table 1).

of X-1 is demonstrated in Figure 1; MCD plus power-law fits to X-1 and X-2 are shown in Figures 2 and 3. The F-test indicates that the improved fits given by the MCD plus power-law model are significant over single-component models at more than the 8σ level of confidence for X-1, and at the 3σ level of confidence for X-2. With the MCD plus power-law model, we measure isotropic luminosities of $L_{X-1} = 2.0^{+1.6}_{-0.3} \times 10^{40}$ erg/s and $L_{X-2} = 6.6^{+1.8}_{-2.0} \times 10^{39}$ erg/s (0.2–10.0 keV, assuming $d = 3.7$ Mpc). If ULX sources are like Galactic black holes and AGN, a corona may be the source of hard X-rays. Extrapolating to the 0.05–100.0 keV band, we find that isotropic luminosities increase by approximately 50%.

We note that some HII regions within NGC 1313 may have low metallicity (50–75% relative to solar abundances; Zaritsky, Kennicutt, & Huchra 1994). The metallicity assumed for absorbing material can impact low-energy spectral measurements. For X-1, our best-fit cool disk plus power-law model represents an improvement at the 5σ level of confidence relative to simple single-component fits with the abundances of O and Fe fixed at $0.25 \times$ solar in the absorption model. With such an absorption model, single-component fits to the spectra of X-2 are acceptable. The low-abundance absorption model is likely inappropriate, however; fits with this model under-estimate the depth of the O edge in the X-1 spectra. The Galactic column density along the line of sight to NGC 1313 is 4.0×10^{20} cm $^{-2}$; our significantly higher measurements may indicate significant neutral absorption within NGC 1313 or intrinsic to the sources (see Table 1).

The most remarkable result of our analysis is that both models require cool accretion disks ($kT \simeq 150$ eV). These temperatures differ from those derived from the spectra of stellar-mass




FIG. 2.— The unfolded MOS-1 and MOS-2 spectra of NGC 1313 X-1. The total spectrum, cool ($kT \simeq 150$ eV) disk component, and power-law components are shown in black, blue, and red, respectively.

black holes in bright states; in such spectra $kT \simeq 1$ keV is typical (e.g., Sobczak et al. 2000) but temperatures can reach to nearly 2 keV in extreme cases. These temperatures also differ considerably from those measured with single-component fits with the MCD model to the spectra of some ULXs, which approach $kT \simeq 2$ keV (e.g., Makishima et al. 2000).

The MCD model is based on the following relationship: $T \propto M^{-1/4}$. This can be used for scaling by writing $(M_{\text{ULX}}/M_{10 M_\odot}) \propto (kT_{10 M_\odot}/kT_{\text{ULX}})^4$. If we assume that $kT = 1$ keV is a typical inner disk color temperature for $10 M_\odot$ black holes for values of L_X/L_{Edd} similar to those being observed in ULX sources, the best-fit MCD color temperatures measured for X-1 and X-2 then imply black hole primaries with masses near $M_{\text{BH}} \sim 2 \times 10^4 M_\odot$. If we assume $kT = 0.5$ keV may be more appropriate for $10 M_\odot$ black holes at high accretion rates, this scaling still suggests masses of $M_{\text{BH}} \sim 1.2 \times 10^3 M_\odot$ for X-1 and X-2.

Shimura & Takahara (1995) derived a hardening factor correction to account for the effects of opacity on the measured disk temperature and inner radius. This correction is very simple: $kT_{\text{corr}} = f^{-1}kT_{\text{obs}}$, and $R_{\text{in,corr}} = \eta f^2 R_{\text{obs}}$ (where $f = 1.7$ is the hardening factor, and $\eta = 0.63$ is valid for $i < 70^\circ$ and accounts for the difference between the innermost radius and the radius of peak temperature; see Sobczak et al. 2000 and Makishima et al. 2000). Merloni, Fabian, & Ross (2000) have found that hardening corrections may not be constant against changes in the physical parameters of the disk.

The normalization of the MCD model allows for more direct estimates of the black hole masses if we assume that $R_{\text{in}} = R_{\text{ISCO}} = 8.85 \text{ km} (M_{\text{BH}}/M_\odot)$ (where R_{ISCO} is the radius of the innermost stable circular orbit), appropriate for Schwarzschild holes. Simply, $M_{\text{BH}} = \eta f^2 (K/\cos[i])^{1/2} \times (d/10 \text{ kpc}) \times (8.85 \text{ km})^{-1}$, where K is the model normalization, and i is the inclination of the system. Using the 90% confidence lower limit MCD normalizations (see Table 1) we find lower-limit masses of $M_{\text{X-1}} \gtrsim 2200 M_\odot$ and $M_{\text{X-2}} \gtrsim 830 M_\odot$ (assuming $i = 0$ and $d = 3.7 \text{ Mpc}$; note that higher values of i and d would increase the minimum mass estimates, as would significant black hole spin).

Defining $L_{\text{Edd}} = 1.3 (M_{\text{BH}}/M_\odot) \times 10^{38} \text{ erg/s}$ (Frank, King, & Raine 2002) and taking L_X in the 0.05–100.0 keV range




FIG. 3.— The unfolded MOS-1 and MOS-2 spectra of NGC 1313 X-2. The total spectrum, cool ($kT \simeq 160$ eV) disk component, and power-law components are shown in black, blue, and red, respectively.

as a better approximation to a bolometric luminosity, we find lower-limit isotropic luminosity masses of $M_{\text{X-1}} \gtrsim 230 M_\odot$ and $M_{\text{X-2}} \gtrsim 70 M_\odot$. These mass estimates are roughly an order of magnitude below those obtained by scaling from the MCD fit parameters; it is possible that X-1 and X-2 are observed at $0.1 L_{\text{Edd}}$, similar to many Galactic black holes. We caution that the mass limits are only as good as the MCD model and our best-fit continuum models.

4. DISCUSSION

We have analyzed the EPIC MOS-1 and MOS-2 spectra of the ULX sources NGC 1313 X-1 and X-2. The spectra statistically require soft and hard components to describe the continuum emission. When the soft components in X-1 and X-2 are fit with models for accretion disks, low disk temperatures are obtained ($kT \simeq 150$ eV). Scaling these temperatures and the normalization of the MCD model suggests that X-1 and X-2 harbor black holes with $M_{\text{BH}} \simeq 10^3 M_\odot$, or higher. Isotropic Eddington luminosity scaling suggests that X-1 and X-2 may harbor black holes with masses on the order of $M_{\text{BH}} \simeq 10^2 M_\odot$. It is possible that we have observed X-1 and X-2 at $L_X \simeq 0.1 L_{\text{Edd}}$; however, it is not clear which scaling method is superior.

Colbert et al. (1995) estimate the radio power at the position of X-1 to be 10^{19} W/Hz at 1.4 GHz, implying an isotropic radio luminosity of 10^{35} erg/s . The radio to X-ray luminosity ratio is then approximately 5×10^{-6} ; this makes it unlikely that relativistic beaming can account for the flux of X-1, because beaming tends to produce flat νF_ν spectra (Fossati et al. 1998). Pakull & Mirioni (2002) have found that X-1 lies in the center of a diffuse H_α nebula with a radius of approximately 240 pc, and a high $[\text{O I}]/\lambda 6300/\text{H}_\alpha$ ratio implying X-ray photoionization. An optical survey of massive young star clusters in nearby galaxies by Larsen (1999) reveals no cluster candidates within approximately 380 pc of our position for X-1. Similarly, Pakull & Mirioni (2002) find that X-2 lies at the center of an H_α nebula with strong $[\text{Si II}]$ and $[\text{O I}]$ lines, implying that X-2 is acting on the local interstellar medium. Larsen (1999) reports no clusters within a few kpc of X-2. These findings suggest that X-1 and X-2 may emit nearly isotropically and illuminate their local nebulae. It is unlikely, then, that models based on anisotropic emission from stellar-mass black holes (e.g., King et al. 2001)

can explain the observed fluxes.

IMBHs may be the endpoints of very massive low-metallicity stars (Heger et al. 2002), or perhaps Population III stars from the era of galaxy formation (Madau & Rees 2001). Miller and Hamilton (2002) have suggested that IMBHs may grow in globular clusters. Ebisuzaki et al. (2001) have proposed that intermediate mass black holes may form in young compact star clusters. If the disk temperature and normalization scalings are correct, our estimates for the masses of X-1 and X-2 are inconsistent with black holes in X-1 and X-2 being the endpoints of low-metallicity or Population III stars; X-1 and X-2 may be the result of growth by mergers.

Though a power-law produced a statistically acceptable fit to the *ASCA* spectrum of X-1 ($\chi^2/\text{d.o.f.} = 244/259$), the values we have measured with the MCD plus power-law model are broadly consistent with those reported by Colbert & Mushotzky (1999) using the same model. We measure a lower disk temperature and harder power-law index in X-2. Makishima et al. (2000) fit two *ASCA* spectra of NGC 1313 X-2 separated by two years with the MCD model, and found $kT = 1.47$ keV and

$kT = 1.07$ keV. Our fits to the *XMM-Newton* spectra of X-2 with only an MCD component are not acceptable, but we measure $kT = 0.91$ keV — consistent with the latter *ASCA* result.

Results from a *Chandra* observation of NGC 5408 X-1 may also reveal a cool disk in a two-component spectrum (Kaaret et al. 2002). We speculate that future observations of ULX sources may also reveal cool accretion disks in some cases. Future theoretical studies of anisotropic accretion flows at high accretion rates may be able to explain such results in terms of stellar-mass black holes.

We wish to thank Albert Kong for his generous communication. We acknowledge useful comments from the referee, Jimmy Irwin, and from Marat Gilfanov, Phil Kaaret, Andrew King, Richard Mushotzky, and, Zheng Zheng. J.M.M. acknowledges support from the NSF through its Astronomy and Astrophysics Fellowship Program. M.C.M. was supported in part by NSF grant AST 0098436. This work is based on observations obtained with *XMM-Newton*, an ESA mission with instruments and contributions directly funded by ESA Member States and the US (NASA).

REFERENCES

- Arnaud, K. A., 1996, *Astronomical Data Analysis Software and Systems V*, eds. G. Jacoby and J. Barnes, p17, ASP Conf. Series vol. 101
- Colbert, E. J. M., & Mushotzky, R. F., 1999, *ApJ*, 519, 89
- Colbert, E. J. M., Petre, R., Schlegel, E. M., & Ryder, S. D., 1995, *ApJ*, 446, 177
- Colbert, E. J., M., & Ptak, A. F., 2002, *ApJS*, 143, 25
- Ebisuzaki, T., et al., 2001, *ApJ*, 562, L19
- Fabbiano, G., 1989, *ARA & A*, 28, 87
- Fabbiano, G., & Trinchieri, G., 1987, *ApJ*, 315, 46
- Fossati, G., Maraschi, L., Celotti, A., Comastri, A., & Ghisellini, G., 1998, *MNRAS*, 299, 433
- Frank, J., King, A. R., & Raine, D. J., 2002, “Accretion Power in Astrophysics”, (Cambridge Univ. Press: Cambridge)
- Froschini, L., et al., 2002, *A & A*, 392, 817
- Heger, A., Woosley, S. E., Fryer, C. L., and Langer, N. L., 2002, to appear in Proc. “From Twilight to Highlight - The Physics of Supernovae” ESO/MPA/MPE Workshop Garching July 2002
- Kaaret, P., Corbel, S., Prestwich, A. H., & Zezas, A., 2002, *Science*, in press
- King, A. R., Davies, M. B., Ward, M. J., Fabbiano, G., & Elvis, M., 2001, *ApJ*, 552, L109
- La Parola, V., Peres, G., Fabbiano, G., Kim, D. W., & Bocchino, F., 2001, *ApJ*, 556, 47
- Larsen, S. S., 1999, *A&AS*, 139, 393
- Madau, P., & Rees, M. J., 2001, *ApJ*, 551, L27
- Makishima, K., et al., 2000, *ApJ*, 535, 632
- Merloni, A. A., Fabian, A. C., & Ross, R. R., 2000, *MNRAS*, 313, 193
- Miller, M. C., & Hamilton, D. P., 2002, *MNRAS*, 330, 232
- Mitsuda, K., et al., 1984, *PASJ*, 36, 741
- Pakull, M. W., & Mirioni, L., 2002, in the proceedings of the symposium “New Visions of the Universe in the XMM-Newton and Chandra Era”, 26–30 November 2001, ESTEC, the Netherlands (astro-ph/0202488)
- Shakura, N. I., & Sunyaev, R. A., 1973, *A & A*, 86, 121
- Shimura, T., & Takahara, F., 1995, *ApJ*, 445, 780
- Sobczak, G. J., et al., 2000, *ApJ*, 544, 993
- Titarchuk, L., 1994, *ApJ*, 434, 313
- Tully, R. B., 1988, *Nearby Galaxies Catalog* (Cambridge: Cambridge Univ. Press)
- Zaritsky, D., Kennicutt, R. C., & Huchra, J. P., 1994, *ApJ*, 420, 87

Spectral Fit Parameters

TABLE 1

Model/Parameter	NGC 1313 X-1	NGC 1313 X-2
power-law		
N_H (10^{21} cm^{-2})	$2.0^{+0.2}_{-0.1}$	$2.7^{+0.3}_{-0.2}$
Γ	1.90 ± 0.05	2.4 ± 0.1
Norm. (10^{-4})	$6.3^{+0.3}_{-0.4}$	$3.9^{+0.4}_{-0.3}$
χ^2/dof	497.7/384	177.7/167
MCD		
N_H (10^{21} cm^{-2})	$0.41^{+0.09}_{-0.08}$	$0.6^{+0.2}_{-0.1}$
kT (keV)	1.41 ± 0.06	$0.91^{+0.06}_{-0.05}$
Norm. (10^{-2})	$3.0^{+0.5}_{-0.4}$	$6.4^{+1.8}_{-1.4}$
χ^2/dof	809.5/384	265.5/167
Bremsstrahlung		
N_H (10^{21} cm^{-2})	1.27 ± 0.09	1.5 ± 0.02
kT (keV)	$5.9^{+0.6}_{-0.5}$	2.6 ± 0.3
Norm. (10^{-4})	6.4 ± 0.2	3.7 ± 0.3
χ^2/dof	573.3/384	205.6/167
Raymond-Smith		
N_H (10^{21} cm^{-2})	1.16 ± 0.08	1.0 ± 0.1
kT (keV)	5.4 ± 0.3	$3.7^{+0.3}_{-0.2}$
Norm. (10^{-3})	1.58 ± 0.05	0.68 ± 0.03
χ^2/dof	749.8/384	252.2/167
CompTT		
N_H (10^{21} cm^{-2})	$1.1^{+0.3}_{-0.2}$	$1.7^{+0.7}_{-0.9}$
kT_{seed} (keV)	0.18 ± 0.01	$0.15^{+0.05}_{-0.04}$
kT_{corona} (keV)	50 [†]	50 [†]
τ	$0.80^{+0.07}_{-0.08}$	$0.37^{+0.07}_{-0.06}$
Norm. (10^{-5})	$2.8^{+0.4}_{-0.3}$	1.7 ± 0.4
χ^2/dof	431.8/383	164.2/166
MCD + power-law		
N_H (10^{21} cm^{-2})	$4.4^{+1.2}_{-0.6}$	3^{+3}_{-1}
kT (keV)	$0.15^{+0.02}_{-0.04}$	$0.16^{+0.16}_{-0.04}$
Norm.	1000^{+2100}_{-200}	180^{+80}_{-60}
Γ	$1.82^{+0.06}_{-0.09}$	$2.3^{+0.2}_{-0.1}$
Norm. (10^{-4})	$6.1^{+0.6}_{-0.9}$	$3.6^{+1.0}_{-0.8}$
χ^2/dof	385.8/382	167.1/165
F ($10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$) ^a	12^{+10}_{-2}	$4.0^{+1.1}_{-1.2}$
$F_{power-law}/F_{total}$	0.33	0.63
$L_{0.2-10}$ ($10^{40} \text{ erg s}^{-1}$) ^b	$2.0^{+1.6}_{-0.3}$	$0.66^{+0.18}_{-0.20}$
$L_{0.05-100}$ ($10^{40} \text{ erg s}^{-1}$) ^b	$3.3^{+4.8}_{-0.3}$	$1.0^{+0.6}_{-0.1}$

NOTE.—Results of fitting simple models to the EPIC MOS spectra of NGC 1313 X-1 and NGC 1313 X-2. The XSPEC model “phabs” was used to measure the equivalent neutral hydrogen column density along the line of sight.

[†] The temperature of the up-scattering corona was not constrained by fits with the XSPEC model “CompTT” and was therefore fixed at a reasonable value.

^a The absorption-corrected or “unabsorbed” flux.

^b The luminosity in the 0.2–10.0 keV or 0.05–100.0 keV range, assuming a distance of 3.7 Mpc.